

# INTEGRAL GROUP RING OF THE MCLAUGHLIN SIMPLE GROUP

V.A. BOVDI, A.B. KONOVALOV

*Dedicated to 60th birthday of Professor V.I. Sushchansky*

**ABSTRACT.** We consider the Zassenhaus conjecture for the normalized unit group of the integral group ring of the McLaughlin sporadic group  $\text{McL}$ . As a consequence, we confirm for this group the Kimmerle's conjecture on prime graphs.

## 1. INTRODUCTION, CONJECTURES AND MAIN RESULTS

Let  $V(\mathbb{Z}G)$  be the normalized unit group of the integral group ring  $\mathbb{Z}G$  of a finite group  $G$ . A long-standing conjecture of H. Zassenhaus (**ZC**) says that every torsion unit  $u \in V(\mathbb{Z}G)$  is conjugate within the rational group algebra  $\mathbb{Q}G$  to an element in  $G$  (see [26]).

For finite simple groups the main tool for the investigation of the Zassenhaus conjecture is the Luthar–Passi method, introduced in [22] to solve it for  $A_5$ . Later M. Hertweck in [17] extended the Luthar–Passi method and applied it for the investigation of the Zassenhaus conjecture for  $PSL(2, p^n)$ . The Luthar–Passi method proved to be useful for groups containing non-trivial normal subgroups as well. For some recent results we refer to [5, 7, 16, 17, 18, 19]. Also, some related properties and some weakened variations of the Zassenhaus conjecture can be found in [1, 23] and [3, 21].

First of all, we need to introduce some notation. By  $\#(G)$  we denote the set of all primes dividing the order of  $G$ . The Gruenberg–Kegel graph (or the prime graph) of  $G$  is the graph  $\pi(G)$  with vertices labeled by the primes in  $\#(G)$  and with an edge from  $p$  to  $q$  if there is an element of order  $pq$  in the group  $G$ . In [21] W. Kimmerle proposed the following weakened variation of the Zassenhaus conjecture:

**(KC)** If  $G$  is a finite group then  $\pi(G) = \pi(V(\mathbb{Z}G))$ .

In particular, in the same paper W. Kimmerle verified that **(KC)** holds for finite Frobenius and solvable groups. We remark that with respect to the so-called  $p$ -version of the Zassenhaus conjecture the investigation of Frobenius groups was completed by M. Hertweck and the first author in [4]. In [6, 7, 8, 9, 10, 12] **(KC)** was confirmed for the Mathieu simple groups  $M_{11}$ ,  $M_{12}$ ,  $M_{22}$ ,  $M_{23}$ ,  $M_{24}$  and the sporadic Janko simple groups  $J_1$ ,  $J_2$  and  $J_3$ .

---

1991 *Mathematics Subject Classification.* Primary 16S34, 20C05, secondary 20D08.

*Key words and phrases.* Zassenhaus conjecture, Kimmerle conjecture, torsion unit, partial augmentation, integral group ring.

The research was supported by OTKA grants No.T 43034, No.K61007.

Here we continue these investigations for the McLaughlin simple group  $\text{McL}$ . Although using the Luthar–Passi method we cannot prove the rational conjugacy for torsion units of  $V(\mathbb{Z}\text{McL})$ , our main result gives a lot of information on partial augmentations of these units. In particular, we confirm the Kimmerle’s conjecture for this group.

Let  $G = \text{McL}$ . It is well known (see [15]) that  $|G| = 2^7 \cdot 3^6 \cdot 5^3 \cdot 7 \cdot 11$  and  $\exp(G) = 2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$ . Let

$$\mathcal{C} = \{C_1, C_{2a}, C_{3a}, C_{3b}, C_{4a}, C_{5a}, C_{5b}, C_{6a}, C_{6b}, C_{7a}, C_{7b}, C_{8a}, C_{9a}, C_{9b}, \\ C_{10a}, C_{11a}, C_{11b}, C_{12a}, C_{14a}, C_{14b}, C_{15a}, C_{15b}, C_{30a}, C_{30b}\}$$

be the collection of all conjugacy classes of  $\text{McL}$ , where the first index denotes the order of the elements of this conjugacy class and  $C_1 = \{1\}$ . Suppose  $u = \sum \alpha_g g \in V(\mathbb{Z}G)$  has finite order  $k$ . Denote by  $\nu_{nt} = \nu_{nt}(u) = \varepsilon_{C_{nt}}(u) = \sum_{g \in C_{nt}} \alpha_g$  the partial augmentation of  $u$  with respect to  $C_{nt}$ . From the Berman–Higman Theorem (see [2] and [25], Ch.5, p.102) one knows that  $\nu_1 = \alpha_1 = 0$  and

$$(1) \quad \sum_{C_{nt} \in \mathcal{C}} \nu_{nt} = 1.$$

Hence, for any character  $\chi$  of  $G$ , we get that  $\chi(u) = \sum \nu_{nt} \chi(h_{nt})$ , where  $h_{nt}$  is a representative of the conjugacy class  $C_{nt}$ .

Our main result is the following

**Theorem 1.** *Let  $G$  denote the McLaughlin simple group  $\text{McL}$ . Let  $u$  be a torsion unit of  $V(\mathbb{Z}G)$  of order  $|u|$ . Denote by  $\mathfrak{P}(u)$  the tuple*

$$(\nu_{2a}, \nu_{3a}, \nu_{3b}, \nu_{4a}, \nu_{5a}, \nu_{5b}, \nu_{6a}, \nu_{6b}, \nu_{7a}, \nu_{7b}, \nu_{8a}, \nu_{9a}, \nu_{9b}, \\ \nu_{10a}, \nu_{11a}, \nu_{11b}, \nu_{12a}, \nu_{14a}, \nu_{14b}, \nu_{15a}, \nu_{15b}, \nu_{30a}, \nu_{30b}) \in \mathbb{Z}^{23}$$

*of partial augmentations of  $u$  in  $V(\mathbb{Z}G)$ . The following properties hold.*

- (i) *There is no elements of orders 21, 22, 33, 35, 55, 77 in  $V(\mathbb{Z}G)$ . Equivalently, if  $|u| \notin \{18, 20, 24, 28, 36, 40, 45, 56, 60, 72, 90, 120, 180, 360\}$ , then  $|u|$  coincides with the order of some element  $g \in G$ .*
- (ii) *If  $|u| = 2$ , then  $u$  is rationally conjugate to some  $g \in G$ .*
- (iii) *If  $|u| = 3$ , then all components of  $\mathfrak{P}(u)$  are zero except possibly  $\nu_{3a}$  and  $\nu_{3b}$ , and the pair  $(\nu_{3a}, \nu_{3b})$  is one of*  

$$\{(-2, 3), (-1, 2), (0, 1), (1, 0)\}.$$
- (iv) *If  $|u| = 5$ , then all components of  $\mathfrak{P}(u)$  are zero except possibly  $\nu_{5a}$  and  $\nu_{5b}$ , and the pair  $(\nu_{5a}, \nu_{5b})$  is one of*  

$$\{(-4, 5), (-3, 4), (-2, 3), (-1, 2), (0, 1), (1, 0)\}.$$
- (v) *If  $|u| = 7$ , then all components of  $\mathfrak{P}(u)$  are zero except possibly  $\nu_{7a}$  and  $\nu_{7b}$ , and the pair  $(\nu_{7a}, \nu_{7b})$  is one of*  

$$\{(\nu_{7a}, \nu_{7b}) \mid -86 \leq \nu_{7a} \leq 87, \quad \nu_{7a} + \nu_{7b} = 1\}.$$
- (vi) *If  $|u| = 11$ , then all components of  $\mathfrak{P}(u)$  are zero except possibly  $\nu_{11a}$  and  $\nu_{11b}$ , and the pair  $(\nu_{11a}, \nu_{11b})$  is one of*  

$$\{(\nu_{11a}, \nu_{11b}) \mid -9 \leq \nu_{11a} \leq 10, \quad \nu_{11a} + \nu_{11b} = 1\}.$$

As an immediate consequence of part (i) of the Theorem we obtain

**Corollary 1.** *If  $G = \text{McL}$  then  $\pi(G) = \pi(V(\mathbb{Z}G))$ .*

## 2. PRELIMINARIES

The following result is a reformulation of the Zassenhaus conjecture in terms of vanishing of partial augmentations of torsion units.

**Proposition 1.** (see [22] and Theorem 2.5 in [24]) *Let  $u \in V(\mathbb{Z}G)$  be of order  $k$ . Then  $u$  is conjugate in  $\mathbb{Q}G$  to an element  $g \in G$  if and only if for each  $d$  dividing  $k$  there is precisely one conjugacy class  $C$  with partial augmentation  $\varepsilon_C(u^d) \neq 0$ .*

The next result now yield that several partial augmentations are zero.

**Proposition 2.** (see [16], Proposition 3.1; [17], Proposition 2.2) *Let  $G$  be a finite group and let  $u$  be a torsion unit in  $V(\mathbb{Z}G)$ . If  $x$  is an element of  $G$  whose  $p$ -part, for some prime  $p$ , has order strictly greater than the order of the  $p$ -part of  $u$ , then  $\varepsilon_x(u) = 0$ .*

The key restriction on partial augmentations is given by the following result that is the cornerstone of the Luthar–Passi method.

**Proposition 3.** (see [22, 17]) *Let either  $p = 0$  or  $p$  a prime divisor of  $|G|$ . Suppose that  $u \in V(\mathbb{Z}G)$  has finite order  $k$  and assume  $k$  and  $p$  are coprime in case  $p \neq 0$ . If  $z$  is a complex primitive  $k$ -th root of unity and  $\chi$  is either a classical character or a  $p$ -Brauer character of  $G$ , then for every integer  $l$  the number*

$$(2) \quad \mu_l(u, \chi, p) = \frac{1}{k} \sum_{d|k} \text{Tr}_{\mathbb{Q}(z^d)/\mathbb{Q}} \{ \chi(u^d) z^{-dl} \}$$

*is a non-negative integer.*

Note that if  $p = 0$ , we will use the notation  $\mu_l(u, \chi, *)$  for  $\mu_l(u, \chi, 0)$ .

Finally, we shall use the well-known bound for orders of torsion units.

**Proposition 4.** (see [13]) *The order of a torsion element  $u \in V(\mathbb{Z}G)$  is a divisor of the exponent of  $G$ .*

## 3. PROOF OF THE THEOREM

Throughout this section we denote  $\text{McL}$  by  $G$ . The character table of  $G$ , as well as the  $p$ -Brauer character tables, which will be denoted by  $\mathfrak{BCI}(p)$  where  $p \in \{2, 3, 5, 7, 11\}$ , can be found using the computational algebra system GAP [15], which derives these data from [14, 20]. For the characters and conjugacy classes we will use throughout the paper the same notation, indexation inclusive, as used in the GAP Character Table Library.

First of all we will investigate units of orders 2, 3, 5, 7 and 11, since the group  $G$  possesses elements of these orders. After this, by Proposition 4, the order of each torsion unit divides the exponent of  $G$ , so to prove the Kimmerle’s conjecture, it remains to consider units of orders 21, 22, 33, 35, 55 and 77. We prove that no units of all these orders do appear in  $V(\mathbb{Z}G)$ .

Now we consider each case separately.

- Let  $u$  be an involution. Using Proposition 2 we obtain that all partial augmentations except one are zero. Thus by Proposition 1 the proof of part (ii) of Theorem 1 is done.
- Let  $u$  be a unit of order 3. By (1) and Proposition 2 we get  $\nu_{3a} + \nu_{3b} = 1$ . Put  $t_1 = 5\nu_{3a} - 4\nu_{3b}$ . By (2) we obtain the system of inequalities

$$\mu_0(u, \chi_2, *) = \frac{1}{3}(-2t_1 + 22) \geq 0; \quad \mu_1(u, \chi_2, *) = \frac{1}{3}(t_1 + 22) \geq 0,$$

from which  $t_1 \in \{-22, -19, -16, -13, -10, -7, -4, -1, 2, 5, 8, 11\}$ . Now for each possible value of  $t_1$  consider the system of linear equations

$$\nu_{3a} + \nu_{3b} = 1, \quad 5\nu_{3a} - 4\nu_{3b} = t_1.$$

Since  $\begin{vmatrix} 1 & 1 \\ 5 & -4 \end{vmatrix} \neq 0$ , this system always has the unique solution. First we select only integer solutions, and then using the condition that all  $\mu_i(u, \chi_j, *)$  are non-negative integers, we obtain only four pairs  $(\nu_{3a}, \nu_{3b})$  listed in part (iii) of Theorem 1.

- Let  $u$  be a unit of order 5. By (1) and Proposition 2 we get  $\nu_{5a} + \nu_{5b} = 1$ . Put  $t_1 = 3\nu_{5a} - 2\nu_{5b}$ . By (2) we obtain the system of inequalities

$$\mu_0(u, \chi_2, *) = \frac{1}{5}(-4t_1 + 22) \geq 0; \quad \mu_1(u, \chi_2, *) = \frac{1}{5}(t_1 + 22) \geq 0,$$

so  $t_1 \in \{-22, -17, -12, -7, -2, 3\}$ . Using the same arguments as in the previous case, we obtain only six pairs  $(\nu_{5a}, \nu_{5b})$  listed in part (iv) of Theorem 1.

- Let  $u$  be a unit of order 7. By (1) and Proposition 2 we get  $\nu_{7a} + \nu_{7b} = 1$ . Put  $t_1 = 4\nu_{7a} - 3\nu_{7b}$ . Using  $\mathfrak{BCT}(3)$  and  $\mathfrak{BCT}(5)$ , by (2) we have

$$\begin{aligned} \mu_3(u, \chi_7, 3) &= \frac{1}{7}(t_1 + 605) \geq 0; & \mu_1(u, \chi_{12}, 5) &= \frac{1}{7}(-t_1 + 3245) \geq 0; \\ \mu_1(u, \chi_7, 3) &= \frac{1}{7}(-3\nu_{7a} + 4\nu_{7b} + 605) \geq 0. \end{aligned}$$

It follows that we have only 174 pairs  $(\nu_{7a}, \nu_{7b})$ , given in part (v) of Theorem 1. Note that using our implementation of the Luthar–Passi method, which we intended to make available in the GAP package LAGUNA [11], we checked that it is not possible to further reduce the number of solutions, and the same remark also applies for the remaining part of the paper.

- Let  $u$  be a unit of order 11. By (1) and Proposition 2 we have  $\nu_{11a} + \nu_{11b} = 1$ . Using  $\mathfrak{BCT}(3)$ , by (2) we obtain the system of inequalities

$$\begin{aligned} \mu_1(u, \chi_3, 3) &= \frac{1}{11}(6\nu_{11a} - 5\nu_{11b} + 104) \geq 0; \\ \mu_2(u, \chi_3, 3) &= \frac{1}{11}(-5\nu_{11a} + 6\nu_{11b} + 104) \geq 0, \end{aligned}$$

that has only that twenty pairs  $(\nu_{11a}, \nu_{11b})$  listed in part (vi) of the Theorem 1.

- Let  $u$  be a unit of order 21. By (1) and Proposition 2 we have

$$\nu_{3a} + \nu_{3b} + \nu_{7a} + \nu_{7b} = 1.$$

Put  $t_1 = 5\nu_{3a} - 4\nu_{3b} - \nu_{7a} - \nu_{7b}$ ,  $t_2 = 5\nu_{3a} + 2\nu_{3b}$  and  $t_3 = 3\nu_{7a} - 4\nu_{7b}$ . Since  $|u^7| = 3$ , for any character  $\chi$  of  $G$  we need to consider four cases, defined by part (iii) of the Theorem. Now we consider each case separately:

Case 1. Let  $\chi(u^7) = \chi(3a)$ . Using (2), we obtain the system of inequalities

$$\mu_3(u, \chi_2, *) = \frac{1}{21}(2t_1 + 11) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{21}(-12t_1 + 18) \geq 0,$$

which has no integral solution.

Case 2. Let  $\chi(u^7) = \chi(3b)$ . Again, using (2), we obtain the system of inequalities

$$\begin{aligned} \mu_0(u, \chi_2, *) &= \frac{1}{21}(-12t_1 + 36) \geq 0; & \mu_7(u, \chi_2, *) &= \frac{1}{21}(6t_1 + 24) \geq 0; \\ \mu_0(u, \chi_3, *) &= \frac{1}{21}(36t_2 + 243) \geq 0; & \mu_7(u, \chi_3, *) &= \frac{1}{21}(-18t_2 + 225) \geq 0; \\ \mu_1(u, \chi_{16}, *) &= \frac{1}{21}(-t_3 + 8386) \geq 0; & \mu_9(u, \chi_{16}, *) &= \frac{1}{21}(2t_3 + 8386) \geq 0; \\ \mu_1(u, \chi_5, *) &= \frac{1}{21}(-13\nu_{3a} + 5\nu_{3b} + 765) \geq 0. \end{aligned}$$

This yields  $t_1 \in \{-4, 3\}$ ,  $t_2 \in \{-5, 2, 9\}$  and  $t_3 \in \{7 + 21k \mid -200 \leq k \leq 399\}$ , but none of possible combinations of  $t_i$ 's gives us any solution.

Case 3. Let  $\chi(u^7) = -2\chi(3a) + 3\chi(3b)$ . Then using (2), we obtain the system

$$\begin{aligned}\mu_1(u, \chi_2, *) &= \frac{1}{21}(-t_1 - 1) \geq 0; & \mu_7(u, \chi_2, *) &= \frac{1}{21}(6t_1 + 6) \geq 0; \\ \mu_0(u, \chi_3, *) &= \frac{1}{21}(36t_2 + 207) \geq 0; & \mu_7(u, \chi_3, *) &= \frac{1}{21}(-18t_2 + 243) \geq 0; \\ \mu_1(u, \chi_{16}, *) &= \frac{1}{21}(-t_3 + 8218) \geq 0; & \mu_9(u, \chi_{16}, *) &= \frac{1}{21}(2t_3 + 8218) \geq 0,\end{aligned}$$

from which  $t_1 = -1$ ,  $t_2 \in \{-4, 3, 10\}$  and  $t_3 \in \{7 + 21k \mid -196 \leq k \leq 391\}$ , and again we have no solution for every combination of  $t_i$ 's.

Case 4. Let  $\chi(u^7) = -\chi(3a) + 2\chi(3b)$ . Using (2), we obtain the system

$$\begin{aligned}\mu_7(u, \chi_2, *) &= \frac{1}{21}(6t_1 + 15) \geq 0; & \mu_0(u, \chi_2, *) &= \frac{1}{21}(-12t_1 + 54) \geq 0; \\ \mu_0(u, \chi_3, *) &= \frac{1}{21}(36t_2 + 225) \geq 0; & \mu_7(u, \chi_3, *) &= \frac{1}{21}(-18t_2 + 234) \geq 0; \\ \mu_9(u, \chi_{16}, *) &= \frac{1}{21}(2t_3 + 8015) \geq 0; & \mu_1(u, \chi_{16}, *) &= \frac{1}{21}(-t_3 + 8015) \geq 0,\end{aligned}$$

so  $t_1 = 1$ ,  $t_2 \in \{-1, 6, 13\}$  and  $t_3 \in \{14 + 21k \mid -191 \leq k \leq 381\}$ , that also gives us no solutions.

- Let  $u$  be a unit of order 22. By (1) and Proposition 2 we have

$$\nu_{2a} + \nu_{11a} + \nu_{11b} = 1.$$

Now by (2) we obtain the system of inequalities

$$\mu_0(u, \chi_2, *) = \frac{1}{22}(60\nu_{2a} + 28) \geq 0; \quad \mu_{11}(u, \chi_2, *) = \frac{1}{22}(-60\nu_{2a} + 16) \geq 0,$$

which has no integral solution.

- Let  $u$  be a unit of order 33. By (1) and Proposition 2 we have

$$\nu_{3a} + \nu_{3b} + \nu_{11a} + \nu_{11b} = 1.$$

Put  $t_1 = 5\nu_{3a} - 4\nu_{3b}$ ,  $t_2 = 5\nu_{3a} + 2\nu_{3b}$  and  $t_3 = 32\nu_{3a} - 4\nu_{3b} - 6\nu_{11a} + 5\nu_{11b}$ . Since  $|u^{11}| = 3$ , for any character  $\chi$  of  $G$  we need to consider four cases, defined by part (iii) of the Theorem.

Case 1. Let  $\chi(u^{11}) = \chi(3a)$ . Then by (2) we obtain the system of inequalities

$$\mu_{11}(u, \chi_2, *) = \frac{1}{33}(10t_1 + 27) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{33}(-20t_1 + 12) \geq 0,$$

that has no integral solution.

Case 2. Let  $\chi(u^{11}) = \chi(3b)$ . Now (2) gives us the system

$$\mu_{11}(u, \chi_2, *) = \frac{1}{33}(10t_1 + 18) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{33}(-20t_1 + 30) \geq 0,$$

which also has no integral solution.

Case 3. Let  $\chi(u^{11}) = -2\chi(3a) + 3\chi(3b)$ . By (2) we obtain that

$$\begin{aligned}\mu_1(u, \chi_2, *) &= \frac{1}{33}(-5\nu_{3a} + 4\nu_{3b}) \geq 0; & \mu_{11}(u, \chi_2, *) &= \frac{1}{33}(50\nu_{3a} - 40\nu_{3b}) \geq 0; \\ \mu_0(u, \chi_3, *) &= \frac{1}{33}(60t_2 + 207) \geq 0; & \mu_{11}(u, \chi_3, *) &= \frac{1}{33}(-30t_2 + 243) \geq 0; \\ \mu_1(u, \chi_7, *) &= \frac{1}{33}(t_3 + 978) \geq 0; & \mu_3(u, \chi_7, *) &= \frac{1}{33}(-2t_3 + 750) \geq 0.\end{aligned}$$

It follows that  $t_1 = 0$ ,  $t_2 = 7$  and  $t_3 \in \{12 + 33k \mid -30 \leq k \leq 11\}$ , and we have no solutions again.

Case 4. Let  $\chi(u^{11}) = -\chi(3a) + 2\chi(3b)$ . By (2) we obtain the system

$$\mu_{11}(u, \chi_2, *) = \frac{1}{33}(10t_1 + 9) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{33}(-20t_1 + 48) \geq 0,$$

which has no integral solution.

- Let  $u$  be a unit of order 35. By (1) and Proposition 2 we have

$$\nu_{5a} + \nu_{5b} + \nu_{7a} + \nu_{7b} = 1.$$

Put  $t_1 = 3\nu_{5a} - 2\nu_{5b} - \nu_{7a} - \nu_{7b}$ ,  $t_2 = 6\nu_{5a} + \nu_{5b}$  and  $t_3 = 6\nu_{5a} + \nu_{5b} + 3\nu_{7a} - 4\nu_{7b}$ . Since  $|u^7| = 5$ , for any character  $\chi$  of  $G$  we need to consider six cases, defined by part (iv) of the Theorem.

Case 1. Let  $\chi(u^7) = \chi(5a)$ . By (2) we obtain the system

$$\mu_5(u, \chi_2, *) = \frac{1}{35}(4t_1 + 9) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{35}(-24t_1 + 16) \geq 0,$$

which has no integral solutions.

Case 2. Let  $\chi(u^7) = \chi(5b)$ . Now the non-compatible inequalities are

$$\mu_7(u, \chi_2, *) = \frac{1}{35}(6t_1 + 26) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{35}(-24t_1 + 36) \geq 0.$$

Case 3. Let  $\chi(u^7) = -2\chi(5a) + 3\chi(5b)$ . By (2) we obtain the system

$$\mu_7(u, \chi_2, *) = \frac{1}{35}(6t_1 + 16) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{35}(-24t_1 + 76) \geq 0,$$

which has no integral solution.

Case 4. Let  $\chi(u^7) = -3\chi(5a) + 4\chi(5b)$ . By (2) we obtain the system

$$\begin{aligned} \mu_0(u, \chi_2, *) &= \frac{1}{35}(-24t_1 + 96) \geq 0; & \mu_7(u, \chi_2, *) &= \frac{1}{35}(6t_1 + 11) \geq 0; \\ \mu_0(u, \chi_3, *) &= \frac{1}{35}(24t_2 + 175) \geq 0; & \mu_7(u, \chi_3, *) &= \frac{1}{35}(-6t_2 + 245) \geq 0; \\ \mu_{15}(u, \chi_{16}, *) &= \frac{1}{35}(4t_3 + 8071) \geq 0; & \mu_1(u, \chi_{16}, *) &= \frac{1}{35}(-t_3 + 8001) \geq 0; \\ \mu_0(u, \chi_2, 3) &= \frac{1}{35}(-96\nu_{5a} + 24\nu_{5b} + 85) \geq 0, \end{aligned}$$

so  $t_1 = 4$ ,  $t_2 \in \{0, 35\}$  and  $t_3 \in \{21 + 35k \mid -58 \leq k \leq 228\}$ , and we have no solutions again.

Case 5. Let  $\chi(u^7) = -4\chi(5a) + 5\chi(5b)$ . Using (2), we obtain

$$\begin{aligned} \mu_7(u, \chi_2, *) &= \frac{1}{35}(6t_1 + 6) \geq 0; & \mu_1(u, \chi_2, *) &= \frac{1}{35}(-t_1 - 1) \geq 0; \\ \mu_0(u, \chi_3, *) &= \frac{1}{35}(24t_2 + 155) \geq 0; & \mu_5(u, \chi_3, *) &= \frac{1}{35}(-4t_2 + 155) \geq 0; \\ \mu_{15}(u, \chi_{16}, *) &= \frac{1}{35}(4t_3 + 8091) \geq 0; & \mu_1(u, \chi_{16}, *) &= \frac{1}{35}(-t_3 + 7996) \geq 0. \end{aligned}$$

Then  $t_1 = 1$ ,  $t_2 \in \{-5, 30\}$  and  $t_3 \in \{16 + 35k \mid -58 \leq k \leq 228\}$ , and we have no solutions as before.

Case 6. Let  $\chi(u^7) = -\chi(5a) + 2\chi(5b)$ . By (2) we have incompatible inequalities

$$\mu_7(u, \chi_2, *) = \frac{1}{35}(6t_1 + 21) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{35}(-24t_1 + 56) \geq 0.$$

- Let  $u$  be a unit of order 55. By (1) and Proposition 2 we have

$$\nu_{5a} + \nu_{5b} + \nu_{11a} + \nu_{11b} = 1.$$

Put  $t_1 = 3\nu_{5a} - 2\nu_{5b}$ ,  $t_2 = 6\nu_{5a} + \nu_{5b}$  and  $t_3 = 4\nu_{5a} - \nu_{5b} + 6\nu_{11a} - 5\nu_{11b}$ . Since  $|u^{11}| = 5$ , for any character  $\chi$  of  $G$  we need to consider six cases, defined by part (iv) of the Theorem.

Case 1. Let  $\chi(u^{11}) = \chi(5a)$ . Then by (2) we obtain incompatible inequalities

$$\mu_5(u, \chi_2, *) = \frac{1}{55}(4t_1 + 10) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{55}(-40t_1 + 10) \geq 0.$$

Case 2. Let  $\chi(u^{11}) = \chi(5b)$ . Using (2) we obtain the system

$$\begin{aligned} \mu_{11}(u, \chi_2, *) &= \frac{1}{55}(10t_1 + 20) \geq 0; & \mu_0(u, \chi_2, *) &= \frac{1}{55}(-40t_1 + 30) \geq 0; \\ \mu_0(u, \chi_3, *) &= \frac{1}{55}(40t_2 + 235) \geq 0; & \mu_{11}(u, \chi_3, *) &= \frac{1}{55}(-10t_2 + 230) \geq 0; \\ \mu_5(u, \chi_7, *) &= \frac{1}{55}(4t_3 + 939) \geq 0; & \mu_1(u, \chi_7, *) &= \frac{1}{55}(-t_3 + 934) \geq 0, \end{aligned}$$

so  $t_1 = -2$ ,  $t_2 \in \{1, 12, 23\}$  and  $t_3 \in \{-1 + 55k \mid -4 \leq k \leq 17\}$ , and in every case we have no integer solution.

Case 3. Let  $\chi(u^{11}) = -2\chi(5a) + 3\chi(5b)$ . By (2) we obtain that

$$\begin{aligned}\mu_{11}(u, \chi_2, *) &= \frac{1}{55}(10t_1 + 10) \geq 0; & \mu_0(u, \chi_2, *) &= \frac{1}{55}(-40t_1 + 70) \geq 0; \\ \mu_0(u, \chi_3, *) &= \frac{1}{55}(40t_2 + 195) \geq 0; & \mu_{11}(u, \chi_3, *) &= \frac{1}{55}(-10t_2 + 240) \geq 0; \\ \mu_5(u, \chi_7, *) &= \frac{1}{55}(4t_3 + 946) \geq 0; & \mu_1(u, \chi_7, *) &= \frac{1}{55}(-t_3 + 891) \geq 0.\end{aligned}$$

From this follows that  $t_1 = -1$ ,  $t_2 \in \{2, 13, 24\}$  and  $t_3 \in \{11 + 55k \mid -4 \leq k \leq 16\}$ , and for every combination of  $t_i$ 's we have no solution.

Case 4. Let  $\chi(u^{11}) = -3\chi(5a) + 4\chi(5b)$ . By (2) we obtain the system

$$\mu_{11}(u, \chi_2, *) = \frac{1}{55}(10t_1 + 5) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{55}(-40t_1 + 90) \geq 0,$$

which has no integral solution.

Case 5. Let  $\chi(u^{11}) = -4\chi(5a) + 5\chi(5b)$ . By (2) we have that

$$\begin{aligned}\mu_{11}(u, \chi_2, *) &= \frac{1}{55}(10t_1) \geq 0; & \mu_1(u, \chi_2, *) &= \frac{1}{55}(-t_1) \geq 0; \\ \mu_0(u, \chi_3, *) &= \frac{1}{55}(40t_2 + 155) \geq 0; & \mu_{11}(u, \chi_3, *) &= \frac{1}{55}(-10t_2 + 250) \geq 0; \\ \mu_5(u, \chi_7, *) &= \frac{1}{55}(4t_3 + 986) \geq 0; & \mu_1(u, \chi_7, *) &= \frac{1}{55}(-t_3 + 881) \geq 0,\end{aligned}$$

so  $t_1 = 0$ ,  $t_2 \in \{3, 14, 25\}$  and  $t_3 \in \{1 + 55k \mid -4 \leq k \leq 16\}$ , and again we have no solutions in every combination of  $t_i$ 's.

Case 6. Let  $\chi(u^{11}) = -\chi(5a) + 2\chi(5b)$ . Using (2) we obtain the system

$$\mu_{11}(u, \chi_2, *) = \frac{1}{55}(10t_1 + 15) \geq 0; \quad \mu_0(u, \chi_2, *) = \frac{1}{55}(-40t_1 + 50) \geq 0,$$

which has no integral solution.

- Let  $u$  be a unit of order 77. By (1) and Proposition 2 we have

$$\nu_{7a} + \nu_{7b} + \nu_{11a} + \nu_{11b} = 1.$$

Then using (2) we obtain the non-compatible system of inequalities

$$\begin{aligned}\mu_0(u, \chi_2, *) &= \frac{1}{77}(60(\nu_{7a} + \nu_{7b}) + 28) \geq 0; \\ \mu_{11}(u, \chi_2, *) &= \frac{1}{77}(-10(\nu_{7a} + \nu_{7b}) + 21) \geq 0.\end{aligned}$$

## REFERENCES

- [1] V. A. Artamonov and A. A. Bovdi. Integral group rings: groups of invertible elements and classical  $K$ -theory. In *Algebra. Topology. Geometry, Vol. 27 (Russian)*, Itogi Nauki i Tekhniki, pages 3–43, 232. Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989.
- [2] S. D. Berman. On the equation  $x^m = 1$  in an integral group ring. *Ukrain. Mat. Ž.*, 7:253–261, 1955.
- [3] F.M. Bleher and W. Kimmerle. On the structure of integral group rings of sporadic groups. *LMS J. Comput. Math.*, 3:274–306 (electronic), 2000.
- [4] V. Bovdi and M. Hertweck. Zassenhaus conjecture for central extensions of  $S_5$ . *J. Group Theory*, pages 1–11, to appear, 2007. (E-print [arXiv:math.RA/0609435v1](#)).
- [5] V. Bovdi, C. Höfert, and W. Kimmerle. On the first Zassenhaus conjecture for integral group rings. *Publ. Math. Debrecen*, 65(3-4):291–303, 2004.
- [6] V. Bovdi, E. Jespers, and A. Konovalov. Torsion units in integral group rings of Janko simple groups. *Preprint*, pages 1–30, submitted, 2007. (E-print [arXiv:math/0608441v3](#)).
- [7] V. Bovdi and A. Konovalov. Integral group ring of the first Mathieu simple group. In *Groups St. Andrews 2005. Vol. I*, volume 339 of *London Math. Soc. Lecture Note Ser.*, pages 237–245. Cambridge Univ. Press, Cambridge, 2007.
- [8] V. Bovdi and A. Konovalov. Integral group ring of the Mathieu simple group  $M_{24}$ . *Preprint*, pages 1–12, submitted, 2007. (E-print [arXiv:0705.1992v1](#)).
- [9] V. Bovdi and A. Konovalov. Integral group ring of the Mathieu simple group  $M_{23}$ . *Comm. Algebra*, pages 1–9, to appear, 2007. (E-print [arXiv:math/0612640v2](#)).

- [10] V. Bovdi, A. Konovalov, and S. Linton. Torsion units in integral group ring of the Mathieu simple group  $M_{22}$ . *Preprint*, pages 1–12, submitted, 2007. (E-print [arXiv:0704.3733v1](https://arxiv.org/abs/0704.3733v1)).
- [11] V. Bovdi, A. Konovalov, R. Rossmanith, and Cs. Schneider. *LAGUNA – Lie AlGebras and UNits of group Algebras, Version 3.4*, 2007. (<http://ukrgap.exponenta.ru/laguna.htm>).
- [12] V. Bovdi, A. Konovalov, and S. Siciliano. Integral group ring of the Mathieu simple group  $M_{12}$ . *Rend. Circ. Mat. Palermo (2)*, 56:125–136, 2007.
- [13] J.A. Cohn and D. Livingstone. On the structure of group algebras. I. *Canad. J. Math.*, 17:583–593, 1965.
- [14] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson. *Atlas of Finite Groups*. Oxford University Press, Eynsham, 1985. Maximal subgroups and ordinary characters for simple groups, With computational assistance from J. G. Thackray.
- [15] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.4.9*, 2006. (<http://www.gap-system.org>).
- [16] M. Hertweck. On the torsion units of some integral group rings. *Algebra Colloq.*, 13(2):329–348, 2006.
- [17] M. Hertweck. Partial augmentations and Brauer character values of torsion units in group rings. *Comm. Algebra*, pages 1–16, to appear, 2007. (E-print [arXiv:math.RA/0612429v2](https://arxiv.org/abs/math.RA/0612429v2)).
- [18] M. Hertweck. Torsion units in integral group rings or certain metabelian groups. *Proc. Edinb. Math. Soc.*, pages 1–22, to appear, 2007.
- [19] C. Höfert and W. Kimmerle. On torsion units of integral group rings of groups of small order. In *Groups, rings and group rings*, volume 248 of *Lect. Notes Pure Appl. Math.*, pages 243–252. Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [20] C. Jansen, K. Lux, R. Parker, and R. Wilson. *An Atlas of Brauer Characters*, volume 11 of *London Mathematical Society Monographs. New Series*. The Clarendon Press Oxford University Press, New York, 1995. Appendix 2 by T. Breuer and S. Norton, Oxford Science Publications.
- [21] W. Kimmerle. On the prime graph of the unit group of integral group rings of finite groups. In *Groups, rings and algebras*, volume 420 of *Contemporary Mathematics*, pages 215–228. AMS, 2006.
- [22] I. S. Luthar and I. B. S. Passi. Zassenhaus conjecture for  $A_5$ . *Proc. Indian Acad. Sci. Math. Sci.*, 99(1):1–5, 1989.
- [23] I. S. Luthar and P. Trama. Zassenhaus conjecture for  $S_5$ . *Comm. Algebra*, 19(8):2353–2362, 1991.
- [24] Z. Marciniak, J. Ritter, S. K. Sehgal, and A. Weiss. Torsion units in integral group rings of some metabelian groups. II. *J. Number Theory*, 25(3):340–352, 1987.
- [25] R. Sandling. Graham Higman’s thesis “Units in group rings”. In *Integral representations and applications (Oberwolfach, 1980)*, volume 882 of *Lecture Notes in Math.*, pages 93–116. Springer, Berlin, 1981.
- [26] H. Zassenhaus. On the torsion units of finite group rings. In *Studies in mathematics (in honor of A. Almeida Costa) (Portuguese)*, pages 119–126. Instituto de Alta Cultura, Lisbon, 1974.

V.A. BOVDI

INSTITUTE OF MATHEMATICS, UNIVERSITY OF DEBRECEN,  
P.O. BOX 12, H-4010 DEBRECEN, HUNGARY  
INSTITUTE OF MATHEMATICS AND INFORMATICS, COLLEGE OF NYÍREGYHÁZA,  
SÓSTÓI ÚT 31/B, H-4410 NYÍREGYHÁZA, HUNGARY  
E-mail address: [vbovdi@math.klte.hu](mailto:vbovdi@math.klte.hu)

A.B. KONOVALOV

SCHOOL OF COMPUTER SCIENCE, UNIVERSITY OF ST ANDREWS,  
JACK COLE BUILDING, NORTH HAUGH, ST ANDREWS, FIFE, KY16 9SX, SCOTLAND  
E-mail address: [konovalov@member.ams.org](mailto:konovalov@member.ams.org)